

***International Conference on Voice Physiology and Biomechanics:  
Modeling Complexity  
Marseille August 18-20, 2004***

***Interactions between basic and clinical research.***

Evaluation of vocal cord nodules

M. Pedersen

MD, FRCS, Dr.M.Sci.

ENT-specialist, consultant phoniatrician

The Medical Center, Voice Unit

Oestergade 17,DK-1100 Copenhagen

e-mail: [m.f.pedersen@dadlnet.dk](mailto:m.f.pedersen@dadlnet.dk) phone: +45 33 15 96 00, fax: +45 33 13 77 05

**Abstract**

## 1. Introduction

After forty years of research in the area of voice function one of the author's batons for the next generation seems to be: evidence.

The set-up of evidence is accepted although not optimal in e.g. the pharmaceutical industry - and cancer research in which field the (chosen) instructor for a protocol on vocal cord nodules was found in the year 2000: J.-P. Pignon in Paris. Erkki Bianco, ENT specialist for singers in Paris, Michala Pedersen, MD and Dr.Phil. in Oxford edited the protocol, and a randomisation was planned. The protocol seemed to be the first for interventions for vocal cord nodules. A Vonod-team for the randomisation was build after Pignon's suggestion, but the team did not succeed to get money, even if the Danish State Health Care Research Foundation in November 2003 approved the present author's application for the study, they have very small funds. Other/several teams and models should be considered. Experiences from a pilot study with it's own protocol will be given.

## 2. Methods

Every time an idea in natural science is cooked down to a hypothesis a protocol (and following trial) is needed to document the evidence. In the protocol the background of the hypothesis is probably the main issue, because all ideas have a historical reference. This is why the statisticians should be feared and handled with care: Evidence is related to the basic background of the relevant field. At best the area of voice research should educate their own statisticians as in other fields.

Inclusion criteria should be handled and thereafter the 0 hypothesis clearly discribed. Types of randomisation and interventions together with types of outcomes and follow-up time must be well documented in relation to the background studies. Thereafter the protocol can be discussed with a relevant statistician who carries out the standardised practical, blinded randomisation and statistical calculations based on the hypothesis and outcome as made by the researchers. A relevant statistical knowledge on the researchers side is necessary now-a-days to argue with statisticians without knowledge of our field. Mostly a two-tailed significance of  $p < 0.01$  is relevant. The Type 1 error (e.g. 1%) and Type II error (e.g. 10 %) gives the power of the analysis (e.g. 90%). The difference % between the intervention arm must be calculated. A good data monitoring and safety group has a high order.

## 3. Materials and Models

A description of the statistically necessary populations/models (e.g. patients, apparatus/s, genes or other scenario) is made. The objectives and outcomes are described based on an intention to treat /to use. The trial design thereafter demands a statistical calculation based on a comparison and randomisation between at least two groups of populations/models. The amount of experiments/ subjects is depending on the outcome and significancy, chosen by the researchers.

The protocol also includes description of ethics, assessment of efficacy and safety, access to documents, data handling and record keeping, finance and reassurance, publication policy, spin-off projects and references. Appendices must include detailed description of procedures including consents

ect. The person/s behind the idea must be aware of the responsibility which cannot be left to the statisticians or the funding.

## 4. Results and Discussion

Only two -recent- evidence based studies (RCTs) have been found that include vocal cord nodules. For laryngo-pharyngeal reflux three evidence based studies (RCTs) have been found. A meta-analysis seems to be possible for the three studies. Examples and references will be given.

*Key words: Voice - research - evidence*

## 1. Introduction

After a generation of research in the area of voice analysis and function where the role of the Vocal Fold Physiology research since 1981 has played a major role, one of the batons for the next generation seems to be: evidence. Till now no internationally famous girls choirs (as evaluated e.g. by the professional music companies) have shown up, even if 850 years of famous boys choirs (as evaluated e.g. by the companies) have presented the one more beautiful CD/DVD than the other. In a thesis it was shown that well trained girls voices before and in puberty had the same qualities as boys [2-5]. An attempt to evaluate male puberty has earlier been made [6] without the acceptance of evidence as the basic tool [7]. The other aspect which lead to the question of evidence was the Cochrane approach of evidence for treatment of vocal cord nodules. Counting the patients and surgeons: 97% of the patients were females and 100% of the surgeons were males, and there was no evidence neither of diagnosis nor treatment (Cochrane 2000) [8].

### 1.1 Background

The set-up of evidence is accepted although not optimal e.g. in the pharmaceutical industry - and cancer research, in which field the instructor was chosen for a protocol on vocal cord nodules in the year 2002: J.-P. Pignon in Paris. Erkki Bianco, ENT specialist for singers in Paris, and Michala Pedersen, MD and Dr.Phil. in Oxford edited the protocol, and a randomisation was planned. Discussions were also made with research fellow Dr. Alice Beranova from Prag. The protocol seemed to be the first for evidence based interventions for vocal cord nodules. A Vonod-team for the randomisation was build after Pignon's suggestion, but the team did not succeed to get money, even if the Danish State Health Care Research Foundation in November 2003 approved the present first author's application for the study, they have very small funds. Other/several teams and models should be considered.

Therefore results from two pilot studies with their own protocols are presented, the first pilot protocol took into account that for dysphonia, medical treatment is necessary of infections, allergies and disorders provoked by the surroundings: a mean score of Voice Related Quality of Life [9] showed, that medical treatment improved the quality of life with a factor of 13.2-20.8, and interestingly a medical voice hygiene group was also better with a factor of 7.1-8.2. [10]. Results are also given from the our second pilot protocol, of treatment of vocal cord nodules, with a comparison between Zagreb and Copenhagen, at both places also to normals. [11] The modified Cape-V test, jitter and shimmer and quality of life were better at both places after medical therapy for among others the infections of the larynges, that seemed to be the overall most common reason for the voice disorder in Copenhagen and in Zagreb as well (table 1-4 Pedersen et al. 2003).

## 2. Methods

Every time an idea in natural science is cooked down to a hypothesis a protocol (and following trial) is needed to document the evidence (clinical trials) [12]. In the protocol for an evidence based study the background of the hypothesis

is probably the main issue, because all ideas have a accepted historical reference even if meta-analysis only can be based on evidence based studies (the Cochrane Handbook, updated version 2003) [14]. This is why the statisticians should be feared and handled with care: Evidence is related to the basic background of the relevant field. At best the area of voice research should educate their own statisticians as in other fields. Inclusion criteria should be handled and thereafter the 0 hypothesis clearly discribed. Types of randomisation and interventions together with types of outcomes and follow-up time must be well documented in relation to the background studies. Thereafter the protocol can be discussed with a relevant statistician who carries out the standardised practical, blinded randomisation and statistical calculations based on the hypothesis and outcome as made by the researchers. A relevant statistical knowledge on the researchers side is necessary now-a-days to argue with statisticians without sufficient knowledge of our field. Mostly a two-tailed significancy of  $p < 0.01$  is relevant. The Type I error (e.g. 1%) and Type II error (e.g. 10 %) and the power of the analysis (e.g. 90%). The difference % between the intervention arm must be calculated. A good data monitoring and safety group has a high order.

## 3. Materials and Models

A discription of the statistically necessary populations/models of instrumentation (e.g. patients, apparatus/s, genes or other scenario) is made. The objectives and outcomes are described based on an intention to treat /to use. The trial design thereafter demands a statistical calculation based on a comparison and randomisation between at least two groups of populations/models. The amount of experiments/ subjects is depending on the outcome and significancy, chosen by the researchers.

The protocol also includes description of ethics, assessment of efficacy and safety, access to documents, datahandling and record keeping, finance and reassurance, publication policy, spin-off projects and references. Appendices must include detailed discription of procedures including consents ect. The person/s behind the idea must be aware of the reponsibility which cannot be left to the statisticians or the funding.

The levels of evidence are (the Cochrane Handbook) : 1a systematic reviews of RCTs, 1b individual RCTs, 1c life or death evidence based controlled prospective trials. 2a systematic reviews of prospective cohort studies with controls, 2b individual prospective cohort studies with controls, 3a prospective systematic cohort studies, 3b individual prospective cohort studies, 4 communication studies. 5 expert opinion.

## 4. Experiments and Results

Only two- recent-evidence based studies (RCTs) have been found apart from the two pilot studies referred to [10,11], that include vocal cord nodules: [15,16], referred to in: Pedersen M and McGlashan J, A Cochrane review update of vocal cord nodules, to be printed. Before that a survey showed that during the later years stroboscopy and objective measure measurements were given, but prospective analyses and works with control groups were not found. [14].

The study by Benninger [15] compared microspot Co2 laser and microdissection for limited vocal fold benign lesions in a blinded prospective study of 37 patients. Time costs was evaluated, statistical analysis involved an evaluation of a design of 80% power to detect a difference of 1.2 SD between the two groups (alpha = 0.05, twosided test) with 12 subjects in each group. A two-sample Student t test was used for continuous

variables and Fischer's Exact test for categorical variables. A secondary analysis was also repeated on patients with complete data. An ANOVA procedure was carried out. In the study by MacKenzie et al. [16] comparing speech therapist treatment with no treatment of non-organic dysphonia including vocal nodules, computer generated random numbers produced in a restricted randomisation form were supplied by an independent worker in a separate department. The intended number of patients to be recruited in the treatment and non-treatment groups was determined by assuming a medium effect size of treatment (0.5 SD units). Greater than 90% power was sought with alpha set at 0.05 (two tailed). The target chosen was 100 patients in each group, which offered 94% power. At the end of the study the power was 82% (70 and 63 patients in the groups). Statistical analysis compared the mean difference in the outcome variables between the groups with and without treatment by speech therapist. Separate analysis was conducted at the end of treatment and at follow-up. An analysis of covariance was used for both the end of treatment and follow-up analysis. Assessment of pathophysiological outcomes needed a categorical approach: a subtraction for each patient was made from baseline (visit one) of visit two and visit three for each feature score. Patients were then assigned to a category (0,1 or 2) according to whether they had improved, deteriorated or stayed the same. A series of Chi-squared tests were made for inter-group comparisons, treatment effect as mean differences at the relevant outcome at visit two and three using an analysis of covariance in SPSS 9/10.

In the prospective randomised pilot study of dysphonia: medical treatment and medical voice hygiene advice approach [10], the statistical analysis was made with the SAS program. The McNemar test was used for the videostroboscopy and t-tests for the quality of life tests. In the results of the phonetograms, the lowest and highest tones in semitones (Hz) and the area in decibels times semitones were compared, one-way analysis of variance was made, geometric means were computed and the difference from before and after were presented as a ratio, after/before. In the prospective randomised pilot study comparing treatment of vocal nodules in Zagreb and Copenhagen [11] the patients were compared with normals of the same age and sex at the same time at both places. The McNemar test was used again for stroboscopies and for the modified Cape-V and quality of life. The jitter and shimmer evaluations were made with t-tests in the SAS program.

The results of the four prospective randomised double blinded studies with relevant follow-up show that for the first one by Benninger there was no difference between the two procedures of surgery. For the second one by MacKenzie et al. there was no effect of speech therapy on the dysphonia in the blinded evaluation. In the third one by Pedersen et al. the quality of life was improved in a group of patients medically treated with relevant immunological, antibiologic and environmental therapy. But the medical voice hygiene advice had a small effect too. In the fourth (pilot) study comparing medical treatment of vocal nodules in Zagreb and in Copenhagen the medical cause of the nodules was infection in both countries, and in both places a significant improvement of the video-stroboscopical pictures, jitter and shimmer in the MDVP, and of a modified Cape-V was found after medical treatment.

For laryngo-pharyngeal reflux as a possible provocator of nodules four evidence based studies (RCTs) have been found that included hoarseness. A protocol for the Cochrane Library (2004) has been made.

The prior conclusion was that no meta-analysis seems to be possible for the three studies that were evidence based [18,19,20]. Still a modified one will show that two of the studies are for and two against acid pump inhibitor treatment.

## 5. Discussion

For singers clinical trials of evidence is the only scientific approach for their pathologies. Christa Ludwig [21] presents the dialogue between the singer, director of orchestra, instructor and teacher. Art develops with new vocal demands [22,23], science develops with new epidemiological understanding [24], scientific theory of music develops [25,26] – and so does the treatment aspect, one of the basics for medical doctors being the analyses of stroboscopy [27]. The next one, possibly is the highspeed video camera of the vocal cords in the clinic, foreseen by the kymographic approach. There is among highly qualified laryngologists a hidden approach of alternative medicine, on the other side many surgeons operate, where they never should, having never understood the demands of evidence (personal communication Jan Olafsson). If the scientists do not accept to make the research of voice evidence based so that every clinician can make statements to the patients of evidence, our field will be handicapped. It is crucial to acknowledge the treatment differences between the sexes, the ages, the environmental provocations [28], infections, allergies ect. before operations are made, because we know that there is no evidence based documentation of the effect of surgery on benign neoplasms of the vocal cords. The quality of life aspects must be taken into account (Voice Handicap Index, SF 36) [29,30], and careful allround advice must be given referring to the scientific status.

## 6. Conclusion

An introduction to four evidence based studies in adults that included vocal nodules has been made, one study showed that the laser could be used for surgical treatment, another that speech therapy in a blinded setting had no evidence, the third one showed effect of medical treatment for hoarseness with reflux, allergy and infection therapy, but also of a medical voice hygiene therapy (with an effect factor of 3:1). The fourth study showed that both in Zagreb and in Copenhagen infections were seen more often in patients with vocal nodules than in others, and medical treatment had an effect in both places measured with the acoustical analysis of jitter and shimmer, the perceptive test of modified Cape-V and quality of life questionnaire. In four evidence based studies of reflux as the cause of dysphonia no connection was found.

## Acknowledgment

This presentation was carried out with the advice from Michala Pedersen, Dr.phil. Oxford, and Erkki Bianco, Doctor of the opera lyrique in Paris.

Statistical advice was made by the East Danish Health Science Forum's Consultant Service

## 7. References:

1. Vocal Fold Physiology, contemporary research and clinical issues. 1 conf. Ed. 1981. Hirano M. Dpt. Otolaryngology Kurume University Medical School. Japan.
2. Pedersen M. Biological development and the normal voice in puberty. Thesis. Oulu. Finland 1997. ISBN 951-42 4592-X (book: Die biologische Entwicklung der Stimme in der Pubertät)
3. Pedersen MF, Møller S, Krabbe S, Bennet P, Svenstrup B. Fundamental voice frequency in female puberty measured with electroglottography during continuous speech as a secondary sex characteristic. A comparison between voice, pubertal stages, oestrogens and androgens. *Int. J Ped Otorhinolaryngol.* 1990;20:17-24.
4. Pedersen MF, Møller S, Krabbe S, Bennett P. Fundamental voice frequency measured by electroglottography during continuous speech. A new exact secondary sex characteristic in boys in puberty. *Int J Ped Otorhinolaryngol.* 1986;11:21-27.
5. Pedersen MF, Møller S, Krabbe S, Munk E, Bennett P. A multivariate statistical analysis of voice phenomena related to puberty in choir boys. *Folia phoniatr* 1985;37:271-78.
6. Hollien H, Green R, Massey K. Longitudinal research on adolescent voice change in males. *J. Acoust. Soc. Am.* 1994; 96: 2646-54.
7. Pocock SJ. *Clinical trials* 1983. Ed. Wiley & sons
8. Pedersen M, McGlashan J. Surgical versus non-surgical interventions for vocal cord nodules. *The Cochrane library.* 2000-01.
9. Hogikyan ND, Wodchis, WP, Terrol JE, Bradford CR, Escfamado RM. *J Voice.* 2000; 14: 378-85.
10. Pedersen M, Beranova A, Møller S. Dysphonia: medical treatment and a medical voice hygiene advice approach. A prospective randomised pilot study. *European Archived of Oto-Rhino-Laryngology.* 2003 electronic version, in print.
11. Pedersen M, Bofac-Stosic N, B. Vocal fold nodules , a prospective study comparing treatment in Zagreb and Copenhagen. Presented at the Union of Phoniatriicians european conf. Toulouse 2003. Abstract. Accepted for EEn (ref in my clinic)
12. McNeill W, Feldman KS. *Continental philosophy.* 1998. Blackwell Publishers Ltd. Oxford. UK
13. *The Cochrane Handbook.* 2004. The Cochrane library.
14. Pedersen M, McGlashan J. Surgical versus non surgical interventions for vocal cord nodules. *The Cochrane Library,* update 2004, to be printed
15. Benninger MS. Microdissection or Microspot Co2 laser for limited vocal fold benign lesions: a prospective randomized trial. *Laryngoscope* 2000; 110: Supplement 1-17.
16. MacKenzie K, Millar A, Wilson J, Sellars C, Deary IJ. Is voice therapy an effective treatment for limited vocal fold benign lesions? A randomised controlled trial. *BMJ* 2001; 323: 658-61.
17. *Multi Dimensional Voice Program.* By the firm Key-Elementrics. United States.
18. Havas T, Huang S, Levy M, Abi-Hanna D, Truskett P, Priestley J, Cox J, Wilson J. Posterior pharyngolaryngitis. Double-blind randomised placebo-controlled trial of proton pump inhibitor therapy. *Aust.J.Otolaryng.* 1999;3:243-46
19. El-Serag HB, Lee P , Buchner A, Inadomi JM, Gavin M, McCarthy DM. Lansoprazole treatment of patients with chronic idiopathic laryngitis: a placebo-controlled trial. *Am J Gastroenterology* 2001;96: 979-83.
20. Noordzij JP, Khidr A, Evans BA, Desper E, Mittal RK, Reibel JF, Levine PA. Evaluation of omeprazole in the treatment of reflux laryngitis: a prospective, placebo-controlled, randomized, double-blind study. *Laryngoscope* 2001;111:2147-51.
21. Ludwig C: *Erinnerungen.*1999.Ed. Henschel Verlag Berlin ISBN 3-89487-337-X
22. Hindemith P. *Elementary training for musicians.* 1946. Ed. Schott and Co. Ltd. London. ISBN 0-901938-16-5.
23. Schoenberg A. *Theory of harmony.* Originally published 1911 by Universal Edition, Vienna. Ed. 1983 Belmont Music Publishers. Faber and Farber Ltd. ISBN 0-571-13078-X.
24. Cochrane A. *Back to the front.* Ed. Bosch FX, Molas R. 2003 Barcelona. ISBN 84- 607 8958-6.
25. Sundin N-G. *Pictures od the history of music (Swedish).* Mirage 1984. ISBN 91-970403-2-0.
26. Sundin N-G. *Aesthetic Criteria for Musical Interpretation.* Thesis. Jyväskylä Finland 1994. ISBN 0075-4633.
27. Schönhärl E. *Die Stroboskopie in der praktischen Laryngologie.* G. Thieme Verlag. Stuttgart. 1960
28. *Occupational Voice: care and cure.* Ed. P.Dejonckere. Kugler publications. The Hague. The Netherlands. 2001. ISBN 90 6299 179 3.
29. *Voice Handicap Index.* In: *Occupational Voice, care and cure.* Ed. P. Dejonckere. Kugler publications. The Hague. The Netherlands. 2001.
30. SF-36. *General quality of life measure, short form.*

**Table 1. Medical groups of included test patients with vocal cord nodules and normal controls. Patients with vocal cord nodules and controls for the pilot study were collected in a period of two months.**

<b>Groups: with vocal nodules from Denmark</b>	<b>10</b>	<b>mean age 30.3y</b>
<b>with vocal nodules from Croatia</b>	<b>6</b>	<b>“ 38.5y</b>
<b>without vocal nodules from Denmark</b>	<b>13</b>	<b>“ 40.0y</b>
<b>without vocal nodules from Croatia</b>	<b>11</b>	<b>“ 39.7y</b>

**Sex – M:0 F:40**

**Inherited illnesses in the family: None**  
**Currently general health state of the subject: all well**

	<b>Denmark</b>		<b>Croatia</b>	
	<b>Tests</b>	<b>Controls</b>	<b>Tests</b>	<b>Controls</b>
<b>Number of persons</b>				
<b>Illnesses in the past of the voice</b>	<b>3</b>	<b>1</b>	<b>4</b>	<b>0</b>
<b>Infection upper airways</b>	<b>5</b>	<b>0</b>	<b>4</b>	<b>0</b>
<b>Allergy upper airways</b>	<b>4</b>	<b>9</b>	<b>0</b>	<b>0</b>
<b>Reflux/helicobakter infect. (GERD and LPRD)</b>	<b>3</b>	<b>3</b>	<b>2</b>	<b>0</b>
<b>Environmental disorders</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>
<b>Others</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>

**Table 2. Voice-related quality of life questionnaire (V-RQOL), results of the prospective pilot study of the new patients before medical treatment and after treatment, one month later, compared with controls.**

Mean score all in percent	Denmark			Croatia		
	before treatment	after	diff%	before treatment	after	diff%
Social emotional domain	64	76.5	12.5	66.5	79.5	13
Physical functioning domain	56.3	64.7	8.4	67	74.3	7.3
Overall voice related quality of life	59.8	74.9	15.1	69.6	76.4	6.8

Statistical calculation with t-test shows a better V-RQOL test in both Denmark and Croatia ( $p < 0.01$ ) after treatment, the score of the physically functioning domain is significantly lower in Copenhagen ( $p < 0.05$ ).

**Table 3. The multidimensional voice program (MDVP) acoustical analysis: mean jitter% and shimmer% of /a/ and /i/ at the first and second examination with one month interval.**

**The first measurement (MDVP) and second measurement of all patients compared with controls. A significant reduction of shimmer% /i/ was found between the first and second examination. For shimmer% /a/ the same tendency was seen but not significantly (p=0.053) For description of sentence jitter%, see text.**

---

	<b>Denmark</b>		<b>Croatia</b>	
1. EXAMINATION	patients	controls	patients	controls
<b>jitter% /a/</b>	<b>2.9</b>	<b>1.1</b>	<b>2.9</b>	<b>1.4</b>
<b>jitter % /i/</b>	<b>2.4</b>	<b>1.8</b>	<b>2.5</b>	<b>1.3</b>
<b>shimmer% /a/</b>	<b>12.3</b>	<b>3.5</b>	<b>11.0</b>	<b>4.9</b>
<b>shimmer % /i/</b>	<b>11.3</b>	<b>4.6</b>	<b>7.7</b>	<b>3.4</b>
<b>sentence jitter%</b>	<b>6.1</b>	<b>4.5</b>	<b>6.0</b>	<b>4.2</b>

---

2. EXAMINATION	patients	controls	patients	controls
<b>jitter% /a/</b>	<b>2.9</b>	<b>1.1</b>	<b>2.0</b>	<b>1.1</b>
<b>jitter % /i/</b>	<b>2.8</b>	<b>1.8</b>	<b>2.0</b>	<b>0.9</b>
<b>shimmer% /a/</b>	<b>7.2</b>	<b>4.3</b>	<b>5.0</b>	<b>2.0</b>
<b>shimmer % /i/</b>	<b>7.2</b>	<b>5.6</b>	<b>5.0</b>	<b>2.3</b>

